

Weekly Crop Situation Report

19.06.2021 to 25.06.2021

Sr#	Institute	Crop	Sowing Area	Pest/Disease/Weeds Infestation	Overall condition of crop	Rainfall mm	Temp.°C	Advisory to farmers	Additional remarks
1	Sugarcane Research Institute, Faisalabad	Sugarcane	776 (000) ha (2nd estimate, Crop reporting services 2020-21)	Stem borer, Whip Smut in plant crop and Weed infestation in neglected fields.	Normal			<ul style="list-style-type: none"> ● Chemical and cultural practices of weed control should be adopted ● In Spring planted sugarcane crop, earthing up should be done ● In September planted apply one bag of Urea and one bag granular/acre ● Irrigate the September and Spring planted sugarcane according to crop requirement and weather forecast ● Regularly visit the crop, if any problem about insect/ pest, and disease will be solved ● Spray of bifenthrin or lamada @ 250-400ml respectively should be sprayed 	Frequent feedback received from the farmers

							<p>in case of attack of black bugs especially on ratoon crop</p> <ul style="list-style-type: none"> ● Apply 30% more fertilizer to the ratoon crop ● Apply Urea fertilizer to the spring planted crop of sugarcane ● Use recommended insecticide to control borer etc attack to the crop 	
2	Vegetable Research Institute, Faisalabad	Spinach		Alternaria Leaf Blight, Aphid and Jassid	Satisfactory		<ul style="list-style-type: none"> ● Judicious use of fertilizers for better seed production as well as better production of fresh crop ● Irrigate the field as per atmospheric condition for better fresh production ● Spray against insects, pests and diseases ● Save the crop from frost in growing area ● Weeds must be eradicated to minimize plant weed competition 	The crop is at seed setting stage hence the fresh crop production is decreasing in yield and quality.
		Tomato		Aphid Jassid, Blight, Grey	Satisfactory		<ul style="list-style-type: none"> ● Judicious use of fertilizers and 	High temperature

				mold.				<ul style="list-style-type: none"> proper irrigation at flowering and fruit development stage ● Spray against insect pests and diseases ● Maintain proper irrigation at flowering and fruit development stages during high temperature days ● Weeds must be eradicated to minimize plant weed competition 	spell during last week hindered the crop productivity and caused reduction in fruit bearing.
	Chilies		Aphid, Thrips, viral infestation	Satisfactory				<ul style="list-style-type: none"> ● Judicious use of fertilizers and proper irrigate the field ● Remove the plastic sheet to manage high temperature effects ● Spray against sucking insects if required ● Keep filed weed free in both tunnels and open field ● Maintain proper irrigation at flowering and fruit development stages during high temperature days 	High temperature spell during last week hindered the crop productivity and caused reduction in fruit bearing.
	Bottle gourd		Red pumpkin beetle, girding	Satisfactory				<ul style="list-style-type: none"> ● Judicious use of fertilizers after each 	High temperature

			weevil and fruit fly				<ul style="list-style-type: none"> ● picking ● Keep the field weed free to remove crop plant and weed competition. ● Maintain proper irrigation at flowering and fruit development stages during high temperature days 	spell during last week hindered the crop productivity and caused reduction in fruit bearing.
	Okra/Lady Finger		Red pumpkin beetle, gray mold, rotening, Aphid & Fungal Diseases.	Satisfactory			<ul style="list-style-type: none"> ● Judicious use of fertilizers for better production ● Fertilizer application after each picking ● Keep the field in weed free condition ● Irrigate the field as per climatic conditions and keep the field in wattar conditions 	Low production due temperature fluctuation and heat waves.
	Bitter gourd		Myrothecium, Leaf minor, Aphid, Jassid, Downy Mildew and viral diseases	Satisfactory			<ul style="list-style-type: none"> ● Judicious use of fertilizers for better production ● Fertilizer application after each picking ● Keep clean the field from weeds ● Irrigate the crop twice in a week for reducing high temperature effects and keep the field 	Low production due temperature fluctuation and heat waves.

								in watar conditions	
		Tinda gourd		Leaf minor, Aphid, Jassid, Downy Mildew and viral diseases	Satisfactory			<ul style="list-style-type: none"> ● Balanced fertigation to boost fruiting spell and crop growth. ● Fertilizer application after each picking. ● Weeds must be eradicated to minimize plant weed competition. ● Irrigate the field twice in a week to mitigate high temperature effects and keep the field in watar conditions 	High temperature spell during last week hindered the crop productivity and caused reduction in fruit bearing.
3	Oilseed Research Institute, Faisalabad	Sunflower		Pests: Nil Disease: Nil Weeds: Nil				<ul style="list-style-type: none"> ● Select loam or heavy loam soil for the cultivation of sesame ● Avoid sandy and water logged soil for its cultivation ● Prepare the soil by ploughing 2 to 3 times followed by planking ● Get seed of approved sesame varieties from registered sale points ● Best sowing time of sesame is from 20 June to 15 July 	

								<ul style="list-style-type: none"> ● Seed rate should be 2 kg/acre ● Add 1 bag of DAP at the time of sowing ● Row to row distance should be 45 cm 	
4	Pulses Research Institute, Faisalabad	Mung	518.02					<p>Kharif Crop:</p> <ul style="list-style-type: none"> ● Prepare soil, arrange input for mung and mash and complete sowing on first monsoon rain when temperature is below 40 degrees <p>Spring sown</p> <p>Mung & Mash:</p> <ul style="list-style-type: none"> ● Eradicate the weeds from fields ● Remained vigilant against insect pest especially, white fly on mash and thrips on mung at this stage. In this case farmers should spray suitable recommended pesticide ● Irrigate the spring sown crop wherever needed ● Apply nitrogen fertilizer on mash crop wherever needed to improve 	
		Mash	11.67						

								the growth	
5	Horticulture Research Institute, Faisalabad	Guava		Infestation of weeds were recorded Remove weeds by ploughing the field	Satisfactory			<ul style="list-style-type: none"> ● Manage mature crop harvesting keeping in view the weather 	
		Date Palm	0.0148	Spray chloproyriphos around the stems which are exposed to red palm weevil and do earthen up				<ul style="list-style-type: none"> ● Arrange the spathes along with fronds to facilitate thinning. ● Start thinning of densely fruit bunches 	
		Ber	0.0135	Arrange lime and copper sulfate for stem pasting against high temperature				<ul style="list-style-type: none"> ● Eradicate weeds from the field ● Do hoeing around grafted plants ● Start annual pruning of bearing plant up to 50% 	
6	Agronomic Research Institute, Faisalabad	Sugarcane			Satisfactory	8.2 mm (Faisalabad) 0.0 mm (Farooqabad, S.Pura) 0.0 mm (Khanewal) 0.5 mm	39.1/24.9°C (Faisalabad) 40.14/25.57 (Farooqabad) 40.14/25.62°C (Khanewal) 41.0/24.8 ° C (Karor, Layyah)	<ul style="list-style-type: none"> ● Irrigate the crop as per the need. Use appropriate insecticide for the control of root borer ● Apply urea to the spring planted crop 	Effective weed control is a prerequisite for ensuring healthier and vigorous crop growth

		Rice				(Karor, Layyah) 18.0 mm (Bahawalpur)	41.0/26.0°C (Bahawalpur)	<ul style="list-style-type: none"> • Complete production technology can be found at http://dai.agripunjab.gov.pk/system/files/RICE%20PLAN%202021-22.pdf. Transplant only the healthy nursery of proper age • Irrigate the crop as per the need • Use appropriate insecticide for the control of sucking insect (Jassid and Thrips) Apply urea to the crop in split dose 	and yield. For any type of assistance/help regarding weed control in all crops, please contact Mr. Muhammad Ashiq (Senior Scientist) of this institute. His contact number is 0300-76 57 249.
		Cotton							
7	Entomological Research Institute, Faisalabad	Sugarcane		Borers Complex 0-1.8% Pyrilla 0-1.7 per leaf Mealybug Nil Whitefly Nil Black bug 0-2.6	Fruit borer and fruit fly are present on guava			<ul style="list-style-type: none"> • Creating awareness among farmers about major insect pests problem and suggested integrated approach for controlling insect pests 	
		Wheat		Crop at maturity					
		Mango		Mango Fruit Fly Nil Mango Hopper 0-1.6 nymph or adult/ branch					

		Citrus		Fruit Fly 0-3.9 % Psylla 0-2.2 per Leafminer 0-4.5% Black Fly 0-1.85 per leaf				
		Guava		Fruit Fly 0-6.85% infestation 0.17/trap/week Fruit Borer 0-0.47 %				
		Vegetables		Brinjal fruit borer 0-6.0% Thrips Below ETL Mites Above ETL Armyworm In patches Cucurbit sucking insects Below ETL Fruit Fly 0-5.95% Jassid 0-0.7 per leaf				
		Rice		Plant Hopper Nil				
		Maize		Stem borer Nil				
8	Fodder Research Institute, Sargodha	Rabi Fodder		Attack of sting bug and lygus bug was observed on Lucerne seed	Good			<ul style="list-style-type: none"> ● Use certified seed for the better production of kharif fodders ● Pest control

				crops.				measures must be taken according to the recommendations of agriculture department	
9	Mango Research Institute, Multan	Mango	0.264	The infestation of fruit fly was recorded on the mango fruits attaining the maturity.	Satisfactory			<ul style="list-style-type: none"> ● A strong message in the form of a slogan “ Batoor Mukkao , Borer Bhagao” was delivered in the farmer’s gathering held during the period under report 	The gradual rise in temperature may result in the early maturity of the mid and late seasonal varieties.
10	Citrus Research Institute, Sargodha	Citrus		<p>Plant Pathology Division Some symptoms of drying of leaves were observed on different citrus varieties. Minor attack of twig blight.</p> <p>Entomology Division Minor infestation of citrus psylla, citrus scale and lemon butterfly was observed. Infestation of leafminer was also observed in nursery</p>	Satisfactory			<ul style="list-style-type: none"> ● Regular pest monitoring should be done ● Apply Imidacloprit + Bifenthrin for the control of all pests @ 2.5 ml/ liter of water ● Install methyl eugenol pheromone traps in the orchards at the rate of 5/acre ● Spray of copper based fungicide like copper hydroxide @ 2.5 gm/ liter of water for citrus canker and Azoxystrobin @ 1 ml/liter of water for fungal diseases is 	

				plantation.				recommended where fruit has been harvested <ul style="list-style-type: none"> ● Application of metalaxyl + mancozeb @ 2 kg/ acre along with thiophenate methyl @ 800 gm / acre for the root borne diseases of citrus is recommended 	
11	PPRI, Faisalabad	Tomato		Grey mold 9%	Satisfactory			Spray the crop with after the cutting of the fodder <ul style="list-style-type: none"> ● Score @ 1 cc/ lit of water ● Amistar top @ 2cc / lit of water ● Sulpher @ 2.5 gm/ lit of water ● Note: Light irrigation during the month of January & February 	
		Cauliflower		Downy mildew 9 %	Satisfactory			Spray the crop thoroughly with <ul style="list-style-type: none"> ● Amistar top @ 2 CC / lit of water ● Score @ 1 CC / lit of water ● Kumulus@ 2gm/ lit of water 	
12	BARI, Chakwal	Groundnut	0.22	Hairy caterpillar attack was observed in some areas,	Satisfactory			<ul style="list-style-type: none"> ● Weeding should be started after three to four weeks to eradicate weeds 	Agricultural Experts should be consulted for

				which was controlled by spraying insecticides. Weeds infestation was also a serious problem, which was eradicated manually and by spraying weedicides.				from groundnut field. Add gypsum @ 200kg per acre at the time of flowering. Use of gypsum can increase pod size and number of pods per plant and also contribute to increase seed quality. Second weeding should be done at flower initiation to facilitate peg penetration for better pod formation	the control of insects & diseases. Farmers can contact on Mobile phone No. 0334562212 5 (Fida Hassan Shah) for the production technology and problems of Groundnut crop.
		Olive		Very mild attack of wooly aphid is being observed at a few orchards.	Satisfactory			<ul style="list-style-type: none"> ● Control the attack of Wooly Aphid by spraying Biphenthrine @4ml/ L of water ● Irrigate new planted olive plants by applying to avoid heat stress ● Avoid stress at fruit hardening stage 	Advisory services are provided to the farmers at the institute as well as on the farms.
13	Arid Zone Research Institute, Bhakkar	Mungbean						<ul style="list-style-type: none"> ● Keenly observe the attack of Thrips and whitefly and effective insecticide should be applied for the control of insect pests 	

								<ul style="list-style-type: none">● Weeds (narrow and broad leaf) should be control by applying recommended weedicide● Irrigation should be applied by keeping the weather forecast	
--	--	--	--	--	--	--	--	--	--